

SCIENTIFIC REPORTS

OPEN

Three Distinct Glutamate Decarboxylase Genes in Vertebrates

Brian P. Grone¹ & Karen P. Maruska²

Received: 07 April 2016

Accepted: 04 July 2016

Published: 27 July 2016

Gamma-aminobutyric acid (GABA) is a widely conserved signaling molecule that in animals has been adapted as a neurotransmitter. GABA is synthesized from the amino acid glutamate by the action of glutamate decarboxylases (GADs). Two vertebrate genes, *GAD1* and *GAD2*, encode distinct GAD proteins: GAD67 and GAD65, respectively. We have identified a third vertebrate GAD gene, *GAD3*. This gene is conserved in fishes as well as tetrapods. We analyzed protein sequence, gene structure, synteny, and phylogenetics to identify *GAD3* as a homolog of *GAD1* and *GAD2*. Interestingly, we found that *GAD3* was lost in the hominid lineage. Because of the importance of GABA as a neurotransmitter, *GAD3* may play important roles in vertebrate nervous systems.

Glutamate decarboxylases (GADs) are essential for the conversion of glutamate to γ -aminobutyric acid (GABA), the predominant inhibitory neurotransmitter in central nervous systems¹. GADs are members of the Group II pyridoxal-5'-phosphate-dependent decarboxylases, which includes decarboxylases that operate on several different substrates². Two GAD proteins found in vertebrate species, GAD67 and GAD65, are encoded by the paralogous genes *GAD1* and *GAD2*, respectively³. While both GADs synthesize GABA and are co-expressed in most vertebrate GABAergic neurons, *GAD1* synthesizes cytoplasmic GABA that is used for extrasynaptic and metabolic purposes and *GAD2* regulates the vesicular pool for release^{4–6}. Nevertheless, *GAD1* and *GAD2* sequences are highly similar to each other, and they share a common intron-exon organization, indicating a common origin⁷.

The evolutionary history of GAD genes is long and diverse. Genes with homology to *GAD* arose before the evolution of eukaryotes⁸. Genes encoding GAD are found, for example, in *Escherichia coli*⁹, *Saccharomyces cerevisiae*¹⁰, *Drosophila melanogaster*¹¹, and *Caenorhabditis elegans*¹². Furthermore, GABA signaling via membrane receptors elicits hyperpolarization in plants as well as mammals, suggesting conserved or convergent roles for the product of GAD enzymatic activity¹³. In most vertebrate species, only two GAD genes have been described. Another gene in the GAD family, GAD-like 1 (*GADL1*) resembles *GAD1* and *GAD2* in sequence, but is expressed in mouse skeletal muscles and kidney rather than in the brain¹⁴. There have also been some hints of greater diversity in vertebrate *GAD* genes.

In addition to the teleost *gad1* and *gad2* genes, a third gene, *gad3*, was found in brain cDNA of the abyssal grenadier (*Coryphaenoides (Nematonurus) armatus*), a benthic teleost fish¹⁵. A similar *gad3* sequence was subsequently identified in the brain cDNA of goldfish (*Carassius auratus*)¹⁶. The sequences of goldfish and abyssal grenadier *gad3* are clearly related to *gad1a*, *gad1b*, and *gad2* sequences, but their evolutionary history remained unknown¹⁶. Furthermore, no *gad3* genes were reported in any species other than grenadier and goldfish. This absence remained an anomaly, since the goldfish (order Cypriniformes), is very distantly related to the abyssal grenadier (order Gadiformes). Recent teleost phylogenies indicate that the Ostariophysians, of which Cypriniformes including goldfish are members, diverged from the Euteleosts, which include the abyssal grenadier, over 250 million years ago¹⁷. Thus, the conservation of a *gad3* gene in these two divergent species suggested that *gad3* was present in an early teleost ancestor. Because the teleost lineage is known to have experienced a whole-genome duplication early in its evolution, one reasonable possibility could therefore have been that *gad3* was a teleost-specific *gad* paralog^{18–21}.

Since the original identification of *gad3* from teleost brain cDNA, many comparative genomic resources have become available. The sequencing of teleost and other vertebrate genomes has been accompanied by the development of databases and software for analyzing the conservation of genes. Sarcopterygii species with sequenced genomes include primitive fishes, e.g. elephant shark²² and coelacanth²³, as well as tetrapods, e.g. chicken²⁴, dog²⁵, human²⁶, Tasmanian devil²⁷, Chinese softshelled turtle²⁸, and *Xenopus*²⁹. Actinopterygii species with sequenced genomes include the spotted gar³⁰ as well as teleosts like fugu¹⁹, medaka³¹, tilapia³², and zebrafish³³.

¹Department of Neurological Surgery, University of California San Francisco, San Francisco, CA, 94143, USA.

²Department of Biological Sciences, Louisiana State University, Baton Rouge, LA, 70803, USA. Correspondence and requests for materials should be addressed to B.P.G. (email: brian.grone@ucsf.edu)

Species	Scientific Name	GAD1	GAD2	GAD3
Chicken	<i>Gallus gallus</i>	ENSGALT00000043162	ENSGALT00000012268	
Burton's mouthbrooder	<i>Astatotilapia burtoni</i>	Gad1a: XM_014332345 Gad1b: XM_014340384	XM_005932121	XM_005950266
Coelacanth	<i>Latimeria chalumnae</i>	ENSLACT00000014577	ENSLACT00000011268	ENSLACT00000005682
Dog	<i>Canis familiaris</i>	ENSCAFT00000049584	ENSCAFT00000006929	ENSCAFT00000000144
Elephant shark	<i>Callorhynchus millii</i>	SINCAMT00000000719	SINCAMT00000011054	SINCAMT00000005039
Fugu	<i>Takifugu rubripes</i>	Gad1a:ENSTRUT00000045798 Gad1b:ENSTRUT00000020549	ENSTRUT00000024751	ENSTRUT00000021119
Abyssal grenadier	<i>Coryphaenoides armatus</i>	AF043268	AF043267	AF043269
Human	<i>Homo sapiens</i>	ENST00000358196	ENST00000376261	ENST00000592477*
Medaka	<i>Oryzias latipes</i>	Gad1a:ENSORLT00000021605 Gad1b:ENSORLT00000011550	ENSORLT00000016248	
Spotted Gar	<i>Lepisosteus oculatus</i>	ENSLOCT00000009532	ENSLOCT00000009370	ENSLOCT00000015874
Tasmanian Devil	<i>Sarcophilus harrisii</i>	ENSSHAT00000013524	ENSSHAT00000015741	ENSSHAT00000004379
Nile tilapia	<i>Oreochromis niloticus</i>	Gad1a:ENSONIT00000011023 Gad1b:ENSONIT00000023558	ENSONIT00000008095	ENSONIT00000008040
Chinese softshelled turtle	<i>Pelodiscus sinensis</i>	ENSPSIT00000002371	ENSPSIT00000019780	ENSPSIT00000019627
Xenopus	<i>Xenopus tropicalis</i>	ENXETT00000040862	ENXETT00000040531	ENXETT00000012900
Zebrafish	<i>Danio rerio</i>	Gad1a:ENDSART00000140425 Gad1b:ENDSART0000003008	ENDSART00000021609	ENDSART00000109561

Table 1. Vertebrate GAD1, GAD2, and GAD3 transcript sequence IDs. *pseudogene transcript sequence.

We used recently generated genomic resources to ask whether *gad3* is present and expressed in species other than the goldfish and abyssal grenadier. Our results revealed a surprisingly broad conservation of *GAD3* in mammals, reptiles, birds, and amphibians, as well as fishes.

Methods

Throughout this paper, we use standard gene nomenclature. For fishes, gene symbols are lowercase and italicized and protein symbols are capitalized. For other vertebrates, human conventions are used: gene symbols in all capitals and italicized, protein symbols in all capitals.

Vertebrate sequence data (Table 1) for *GAD1*, *GAD2*, and *GAD3* homolog transcripts were downloaded from Ensembl genomes for the following species: chicken (*Gallus gallus*), coelacanth, fugu, human, medaka, spotted gar, Tasmanian devil, tilapia, Chinese softshell turtle, xenopus, zebrafish³⁴. Transcript DNA sequences for elephant shark were retrieved from the elephant shark Ensembl server. Transcript cDNA sequences for grenadier were retrieved from NCBI^{15,35}. The spotted gar genome shares extensive similarity with both tetrapod and teleost genomes, so we chose to focus on this species for sequence alignment, phylogenetics, and intron/exon structure comparisons³⁰. Additionally, we obtained sequences for fruitfly (*Drosophila melanogaster*) *GAD1*: NM_079190, sea urchin (*Strongylocentrotus purpuratus*) *GAD*: XM_779763, amphioxus (*Branchiostoma floridae*) *GAD*: XP_002592141, and tunicate (*Ciona intestinalis*) *GAD*: ENSCINT00000004013.

In addition to the species listed in Table 1, we identified several other tetrapod species with *GAD3* genes. These included: Orangutan: ENSPPYG00000009199, Rhesus: ENSMMUG00000001554, Rabbit: ENSOCUG000000022124, Horse: ENSECAG00000009017, Platypus: ENSOANG00000002106, Lizard: ENSACAG00000008555.

Sequence Alignment. Both DNA and amino acid sequences were aligned using MAFFT v7.017^{36,37} (by translation alignment for CDS sequences); algorithm E-INS-I; scoring matrix: BLOSUM62; gap open penalty: 1.53; offset value: 0.

Model Testing. MEGA 6 software was used to compare 24 DNA evolution models for the aligned *GAD* CDS sequences³⁸. A generalized time-reversible plus gamma (GTR + G + I) model had the lowest BIC score (Bayesian Information Criterion) and AICc value (Akaike Information Criterion, corrected), so it was used for subsequent phylogenetic analyses. In this model, non-uniformity of evolutionary rates among sites is modeled by estimating a discrete Gamma distribution (+G) of rates and by assuming that certain sites are evolutionarily invariable (+I).

Bayesian Phylogenetic Inference. MrBayes 3.2.6³⁹ was used to infer phylogenetic relationships between *GAD* homologs based on aligned nucleotide CDS sequences, and was accessed via the CIPRES web portal⁴⁰. In MrBayes, the GTR + G + I model of evolution was used; with default settings except for the following specified parameters: nruns(number of runs) = 2; ngen(number of generations) = 1000000; samplefreq = 500; nchain (number of chains) = 8; temp(chain heating temperature) = 0.1; savebrlens = yes; burninfrac(fraction of initial generations discarded) = 0.25.

Diagnostics of the MCMC sampling were carried out using Tracer v1.6 (<http://tree.bio.ed.ac.uk/software/tracer/>). The effective sample size (ESS) for each parameter was >300 for each run, allowing adequate sampling of the Markov chain.

The tree file generated using MrBayes was visualized using FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Synteny. Gene synteny for *GAD3* genes was compared to the syntenic region near *GAD3* using Genomicus⁴¹. Orangutan was used as a reference species for cross-species synteny and protein similarity to highlight




Figure 1. Alignment of predicted protein sequences translated from spotted gar (*Lepisosteus oculatus*) GAD genes. Black: amino acids similar in all three sequences; Gray: similar to one corresponding residue; White: not similar to either corresponding residue. Sequence similarity was calculated using BLOSUM62 matrix with threshold = 1.

conservation of synteny and GAD3 protein sequence in vertebrates despite the absence of GAD3 in some hominids. For comparing primate synteny, simiiformes (last common ancestor of simians) was used as the reference taxon. Synteny data, protein similarity, and species images were downloaded from Genomicus.

Results

We found previously uncharacterized GAD3 genes in many vertebrate genomes, including diverse fishes and tetrapods. A teleost *gad3* transcript was found in a transcriptome library generated from testis tissue from *Astatotilapia burtoni*: (>comp56037_c0_seq1_indA_testis). Zebrafish *gad3* has been previously referred to with the identifier *zgc:163121*. Interestingly, GAD3 had already been annotated in the *Xenopus tropicalis* genome as GAD1.2. It appears, however, that it has not yet been studied in *Xenopus*.

Sequence Similarity. Gad3 predicted protein sequence from spotted gar (*Lepisosteus oculatus*) is more similar to Gad1 (Pairwise Identity: 60.5%) than to Gad2 (Pairwise Identity: 53.9%) (Fig. 1). Gad1 and Gad2 share 67.1% pairwise identity. The N-terminal domain, which is quite variable between Gad1 and Gad2, is truncated and highly divergent in Gad3. The N-terminal 92 amino acids (aa) of Gad1 align to the N-terminal 84 aa of Gad2. Gad3 has 42 aa aligning in this range, only 27 of which align to Gad1 and Gad2 sequence (with a 15 aa gap). These 27 aa of Gad3 have: 23.1% pairwise identity with Gad1, 11.5% pairwise identity with Gad2.

Phylogeny. Pyridoxal 5'-phosphate (PLP)-dependent decarboxylase genes include Glutamate Decarboxylase-Like 1 (GADL1), Cysteine Sulfinic Acid Decarboxylase (CSAD), and histidine decarboxylase (HDC), in addition to GADs. Therefore, we tested the phylogenetic relationship of GAD3 to other genes in this group, using HDC as an outgroup for GAD, GADL1, and CSAD genes⁴². A neighbor-joining tree of aligned predicted amino acid sequences from the spotted gar (*Lepisosteus oculatus*) place *gad3* most closely related to the *gad1/gad2* clade (Fig. 2).

A phylogenetic tree of vertebrate GAD1, GAD2, and GAD3 nucleotide coding sequences was generated using MrBayes (Fig. 3). In insects, GAD1 is the single homolog of vertebrate GAD genes (insect GAD2 is homologous to vertebrate CSAD and GADL1). Therefore we chose *Drosophila melanogaster* GAD1 as the outgroup for the vertebrate and deuterostome GAD genes.

Exon-intron Structure. Spotted gar *gad1* and *gad2* each have 16 exons (Fig. 4). Spotted gar *gad3* has 17 exons. While both *gad1* and *gad2* have coding sequence beginning in exon 1, *gad3* coding sequence (CDS) begins in the second exon (exon 2). The predicted coding sequence of *gad3* has a gap (does not align) with the 5' CDS sequence found in *gad1* and *gad2* exon 1, exon 2, and part of exon 3. The 3' portion of the *gad3* CDS is included on exon 17, while *gad1* and *gad2* stop codons are found in exon 16.

Aside from these differences, *gad3* exon structure is largely similar to *gad1* and *gad2*. All of the exon junctions from exon 3 to exon 16 are in identical locations for all three *gad* genes. In our alignment of the three *gad* genes, the only gaps introduced in *gad3* are located in exon 2 and exon 17.

Synteny. In spotted gar, *gad1* and *gad2* are located adjacent to the myosin genes *myo3b* and *myo3a*, respectively. Similarly, in humans GAD1 is located near MYO3B on chromosome 2, and GAD2 is located adjacent to




Figure 2. GAD3 is closely related to GAD1 and GAD2, and more distantly related to other members of the PLP-dependent decarboxylase gene family. A phylogenetic tree of aligned PLP-dependent decarboxylase amino acid sequences from the spotted gar was generated using neighbor-joining and 2000 bootstrap iterations. Percent bootstrap support for nodes are shown. The scale bar (bottom) indicates substitutions per site.




Figure 3. Phylogenetic tree of GAD1, GAD2, and GAD3 nucleotide sequences. This consensus tree was generated using MrBayes with *Drosophila melanogaster* GAD1 as the outgroup. Nodes are labeled with posterior probabilities. Distinct gene lineages are indicated by colors. The scale bar (bottom) indicates substitutions per site.

MYO3A on chromosome 10. On the other hand, *gad3* is not located near a myosin gene in the genome of spotted gar. The genes located adjacent to spotted gar *gad3* are *mc4r* and *cdh20*. This syntenic block of genes is conserved




Figure 4. Shared exon-intron structure of GAD genes from spotted gar (*Lepisosteus oculatus*). (A) Maps of the three *gad* genes showing 16 exons in *gad1* and *gad2*, and 17 exons in *gad3*. The direction of transcription is from left to right. Genomic distance spanned and strand on which the gene is located are indicated for each gene. (B) Aligned spotted gar *gad1*, *gad2*, and *gad3* CDS regions annotated with position of exons. Bases colored black indicate disagreements with the consensus sequence of the three *gad* genes.

across many vertebrate species, and represents the inferred ancestral state of the bony vertebrates (euteleostomi) (Fig. 5).

GAD3 Conservation and Gene Loss. GAD3 predicted protein sequence is highly conserved in diverse vertebrate genomes (Fig. 6). Yet primates appear to have experienced varying degrees of gene loss at the GAD3 locus (Fig. 7). We identified a human transcript (ENST00000592477) with homology to GAD3 (Table 1), derived from a pseudogene located in the human genome in the conserved GAD3 syntenic position between *MC4R* and *CDH20* (Fig. 7). Similarly, gorilla (*Gorilla gorilla*) GAD3 is annotated as a pseudogene in Ensembl (ENSGGOG00000027455). Although macaque (*Macaca mulatta*) and orangutan (*Pongo pygmaeus*) predicted GAD3 protein sequences share relatively high pairwise identity (84.5%), the GAD3 genes in these two species appear to have large insertions (or deletions) in their predicted coding sequence.

Discussion

We identified a novel glutamate decarboxylase homolog, GAD3, found in many vertebrate genomes. We provide phylogenetic and intron/exon structural evidence that GAD3 is an ancient paralog of GAD1 and GAD2. The conserved chromosomal synteny of GAD3 in vertebrates supports an ancient origin for this gene. Surprisingly, GAD3 was lost in the hominid lineage. Taken together, the phylogenetic analyses, comparisons of gene structure, and synteny data suggest that GAD3 arose via gene duplication of a protovertebrate GAD homolog, likely before the duplication of another paralog which gave rise to GAD1 and GAD2.

GAD3 Evolution. Although our data do not rule out the possibility of a local duplication that gave rise to GAD3, they are consistent with an origin of GAD3 in an early vertebrate via whole-genome duplication. Whole-genome duplication is thought to have played a major role in early vertebrate evolution^{43–45}. Following genome duplication, these duplicated gene pairs (ohnologs) experienced a range of outcomes including non-functionalization, sub-functionalization, and neo-functionalization^{46,47}. Sub-functionalization may happen via protein changes⁴⁸ or via regulatory element loss⁴⁹ in which ancestral expression domains are differentially lost in different genes⁵⁰. For example, recent evidence indicates that duplication of a corticotropin-releasing hormone




Figure 5. *GAD3* genes are found in a conserved syntenic region in vertebrates. Compared to orangutan *GAD3* for reference, other species including representatives of mammals, other tetrapods, and Euteleostomi (bony vertebrates) in general have similar chromosomal positions of *GAD3* genes. The central green pentagons (surrounded by a vertical rectangle) represent the *GAD3* genes. For each species, *GAD3* and 10 flanking genes on each side are represented by colored pentagons. The pentagons point in the direction of transcription, and each color identifies a set of orthologous genes. To the right of the figure, both species and chromosome are indicated for each *GAD3* ortholog. Figure modified from Genomicus PhyloView output⁴¹.

(*CRH*) gene in an early vertebrate led to a broadly expressed *CRH1* and a *CRH2* with expression restricted to a single hindbrain nucleus⁵¹. Like our recent analyses of *CRH* genes, the discovery of *GAD3* as a conserved vertebrate gene relied on freely available genomic resources, pointing to the likelihood that many gene families have unannotated homologs remaining to be found in sequenced genomes^{51,52}.

GAD3 Function. *GAD3* is phylogenetically closer to *GAD1* and *GAD2* than to *GADL1*, but nonetheless it is possible that its enzymatic functions differ from those of *GAD1* and *GAD2*. The absence of much of the N-terminal region, which regulates intracellular localization of *GAD1* and *GAD2* proteins, suggests that *GAD3* protein may have different localization⁵.

Little is known regarding the function of *GADL1* enzyme, though polymorphisms are linked to differential response to lithium treatment for bipolar disorder⁵³. Mammalian *GADL1* does not appear to have glutamate decarboxylase activity, despite its name. Instead, it catalyzes the decarboxylation of aspartate, cysteine sulfinic acid, and cysteic acid to β -alanine, hypotaurine, and taurine, respectively¹⁴. Recently, *GADL1* and *CSAD* were found to have preference for cysteine sulfinic acid as a substrate⁴². Future studies of *GAD3* biochemical substrates will be necessary to address the possibility of substrates other than glutamate.

Intriguingly, zebrafish *gad3* (referred to as *zgc163121*) mRNA expression was significantly downregulated by treatment with dexamethasone, a glucocorticoid agonist, in 25hpf larval zebrafish, as measured by microarray and qPCR⁵⁴. In the deep-sea fish in which *gad3* was first described, the armed grenadier, *Coryphaenoides (Nematonurus) armatus*, *gad2* mRNA levels were found to be expressed in the brain in a sexually dimorphic manner, i.e. higher in male hypothalamus than in female, but no differences were found in *gad3* levels³⁵. In the goldfish, however, *gad3* mRNA levels in the telencephalon were highest in sexually mature fish of both sexes during the breeding period⁵⁵. Since the specific role of *gad3* is unknown in any taxa, the full range of factors that regulate *gad3* expression in the brain, and potentially elsewhere, awaits further investigation.

Loss in hominids. The loss of *GAD3* in both chimpanzees and humans appears to have been preceded by changes to *GAD3* sequences in other hominids. Predicted gorilla, orangutan, and gibbon *GAD3* transcripts appear to be truncated relative to fish *gad3* sequence, but it may be that not all the exons in these sequences are fully annotated in Ensembl. Glutamate metabolic pathways appear to have been under positive selection in hominids, as seen for example in the origin of glutamate dehydrogenase 2 (*GLUD2*) by retroposition of *GLUD1*⁵⁶.

Gene losses have played major roles in human evolution⁵⁷. For example, loss of L-gulonolactone oxidase (*GULO*) makes humans and other Haplorhini susceptible to scurvy, a vitamin C deficiency. Despite conferring this disadvantage, *GULO* gene loss has occurred in multiple mammalian lineages, including guinea pigs⁵⁸ and some bats⁵⁹.





Figure 6. *GAD3* protein sequence is highly conserved across Euteleostomi. Predicted protein sequences of *GAD3* orthologs were compared using Genomicus⁴¹. The degree of similarity to the reference sequence (orangutan, in green) is indicated by the color of the block, according to the scale shown at bottom.




Figure 7. *GAD3* is lost in hominids. The central green pentagons (surrounded by a vertical rectangle) represent the *GAD3* genes. Human and chimpanzee genomes have no functional gene at the location corresponding to *GAD3* in other primates. Humans have a pseudogene at this location. A thick blue line between two genes indicates a “gap”, i.e. a gene lost relative to the ancestral Simiiformes genome. A thin blue line between two genes indicates a “break” in the continuity of the alignment, i.e. a gene added or lost relative to the ancestral Simiiformes genome. Figure modified from Genomicus AlignView⁴¹, which relies on data and images from Ensembl³⁴.

Hominids are not the only lineage that has lost *GAD3*. Rodent genomes, including mice, rats, and squirrels also appear to be missing *GAD3* homologs. The absence of *GAD3* in both humans and mice likely explains why this gene was not discovered sooner, since many investigators choose to focus on these two species.

References

1. Roberts, E. Gamma-aminobutyric acid. *Scholarpedia* 2, doi: 10.4249/scholarpedia.3356 (2007).
2. Sandmeier, E., Hale, T. I. & Christen, P. Multiple evolutionary origin of pyridoxal-5 -phosphate-dependent amino acid decarboxylases. *European journal of biochemistry/FEBS* 221 (1994).
3. Erlander, M. G., Tillakaratne, N. J., Feldblum, S., Patel, N. & Tobin, A. J. Two genes encode distinct glutamate decarboxylases. *Neuron* 7 (1991).
4. Kaufman, D. L., Houser, C. R. & Tobin, A. J. Two forms of the gamma-aminobutyric acid synthetic enzyme glutamate decarboxylase have distinct intraneuronal distributions and cofactor interactions. *J Neurochem* 56 (1991).

5. Soghomonian, J. J. & Martin, D. L. Two isoforms of glutamate decarboxylase: why? *Trends Pharmacol Sci* **19**, 500–505 (1998).
6. Tian, N. *et al.* The role of the synthetic enzyme GAD65 in the control of neuronal gamma-aminobutyric acid release. *Proc Natl Acad Sci USA* **96** (1999).
7. Bu, D. F. & Tobin, A. J. The exon-intron organization of the genes (GAD1 and GAD2) encoding two human glutamate decarboxylases (GAD67 and GAD65) suggests that they derive from a common ancestral GAD. *Genomics* **21**, doi: 10.1006/geno.1994.1246 (1994).
8. Jackson, F. R. Prokaryotic and eukaryotic pyridoxal-dependent decarboxylases are homologous. *Journal of molecular evolution* **31** (1990).
9. Smith, D. K., Kassam, T., Singh, B. & Elliott, J. F. *Escherichia coli* has two homologous glutamate decarboxylase genes that map to distinct loci. *J Bacteriol* **174** (1992).
10. Coleman, S. T., Fang, T. K., Rovinsky, S. A., Turano, F. J. & Moye-Rowley, W. S. Expression of a glutamate decarboxylase homologue is required for normal oxidative stress tolerance in *Saccharomyces cerevisiae*. *J Biol Chem* **276**, doi: 10.1074/jbc.M007103200 (2001).
11. Jackson, F. R., Newby, L. M. & Kulkarni, S. J. Drosophila GABAergic systems: sequence and expression of glutamic acid decarboxylase. *Journal of Neurochemistry* **54** (1990).
12. Jin, Y., Jorgensen, E., Hartwig, E. & Horvitz, H. R. The *Caenorhabditis elegans* gene *unc-25* encodes glutamic acid decarboxylase and is required for synaptic transmission but not synaptic development. *The Journal of neuroscience: the official journal of the Society for Neuroscience* **19** (1999).
13. Ramesh, S. A. *et al.* GABA signalling modulates plant growth by directly regulating the activity of plant-specific anion transporters. *Nat Commun* **6**, doi: 10.1038/ncomms8879 (2015).
14. Liu, P. *et al.* Role of glutamate decarboxylase-like protein 1 (GADL1) in taurine biosynthesis. *The Journal of biological chemistry* **287**, doi: 10.1074/jbc.M112.393728 (2012).
15. Bosma, P. T. *et al.* Multiplicity of glutamic acid decarboxylases (GAD) in vertebrates: molecular phylogeny and evidence for a new GAD paralog. *Molecular biology and evolution* **16** (1999).
16. Lariviere, K. *et al.* GAD(65) and GAD(67) isoforms of the glutamic acid decarboxylase gene originated before the divergence of cartilaginous fishes. *Molecular biology and evolution* **19** (2002).
17. Near, T. J. *et al.* Resolution of ray-finned fish phylogeny and timing of diversification. *P Natl Acad Sci USA* **109**, doi: 10.1073/pnas.1206625109 (2012).
18. Amores, A., Catchen, J., Ferrara, A., Fontenot, Q. & Postlethwait, J. H. Genome evolution and meiotic maps by massively parallel DNA sequencing: spotted gar, an outgroup for the teleost genome duplication. *Genetics* **188**, doi: 10.1534/genetics.111.127324 (2011).
19. Christoffels, A. *et al.* Fugu genome analysis provides evidence for a whole-genome duplication early during the evolution of ray-finned fishes. *Mol Biol Evol* **21**, doi: 10.1093/molbev/msh114 (2004).
20. Hoegg, S., Brinkmann, H., Taylor, J. S. & Meyer, A. Phylogenetic timing of the fish-specific genome duplication correlates with the diversification of teleost fish. *J Mol Evol* **59**, doi: 10.1007/s00239-004-2613-z (2004).
21. Jaillon, O. *et al.* Genome duplication in the teleost fish *Tetraodon nigroviridis* reveals the early vertebrate proto-karyotype. *Nature* **431**, doi: nature03025 (2004).
22. Venkatesh, B. *et al.* Elephant shark genome provides unique insights into gnathostome evolution. *Nature* **505**, doi: 10.1038/nature12826 (2014).
23. Amemiya, C. T. *et al.* The African coelacanth genome provides insights into tetrapod evolution. *Nature* **496**, doi: 10.1038/nature12027 (2013).
24. Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* **432**, doi: 10.1038/nature03154 (2004).
25. Kirkness, E. F. *et al.* The dog genome: survey sequencing and comparative analysis. *Science* **301**, doi: 10.1126/science.1086432 (2003).
26. Venter, J. C. *et al.* The sequence of the human genome. *Science* **291**, doi: 10.1126/science.1058040 (2001).
27. Murchison, E. P. *et al.* Genome sequencing and analysis of the Tasmanian devil and its transmissible cancer. *Cell* **148**, doi: 10.1016/j.cell.2011.11.065 (2012).
28. Wang, Z. *et al.* The draft genomes of soft-shell turtle and green sea turtle yield insights into the development and evolution of the turtle-specific body plan. *Nature genetics* **45**, doi: 10.1038/ng.2615 (2013).
29. Hellsten, U. *et al.* The genome of the Western clawed frog *Xenopus tropicalis*. *Science* **328**, doi: 10.1126/science.1183670 (2010).
30. Braasch, I. *et al.* The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. *Nat Genet* **48**, doi: 10.1038/ng.3526 (2016).
31. Kasahara, M. *et al.* The medaka draft genome and insights into vertebrate genome evolution. *Nature* **447**, doi: 10.1038/nature05846 (2007).
32. Brawand, D. *et al.* The genomic substrate for adaptive radiation in African cichlid fish. *Nature* **513**, doi: 10.1038/nature13726 (2014).
33. Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to the human genome. *Nature* **496**, doi: 10.1038/nature12111 (2013).
34. Flicek, P. *et al.* Ensembl 2014. *Nucleic acids research* **42**, doi: 10.1093/nar/gkt1196 (2014).
35. Trudeau, V. L., Bosma, P. T., Collins, M., Priede, I. G. & Docherty, K. Sexually dimorphic expression of glutamate decarboxylase mRNA in the hypothalamus of the deep sea armed grenadier, *Coryphaenoides (Nematonurus) armatus*. *Brain, behavior and evolution* **56**, doi: 47210 (2000).
36. Katoh, K., Kuma, K., Toh, H. & Miyata, T. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res* **33**, doi: 10.1093/nar/gki198 (2005).
37. Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30** (2002).
38. Tamura, K., Stecher, G., Peterson, D., Filipowski, A. & Kumar, S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* **30**, doi: 10.1093/molbev/mst197 (2013).
39. Ronquist, F. *et al.* MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* **61**, doi: 10.1093/sysbio/sys029 (2012).
40. Miller, M. A., Pfeiffer, W. & Schwartz, T. In Proceedings of the Gateway Computing Environments Workshop (GCE) (2010).
41. Louis, A., Muffato, M. & Roest Crollius, H. Genomicus: five genome browsers for comparative genomics in eukaryota. *Nucleic acids research* **41**, doi: 10.1093/nar/gks1156 (2013).
42. Winge, I. *et al.* Mammalian CSAD and GADL1 have distinct biochemical properties and patterns of brain expression. *Neurochem Int* **90**, doi: 10.1016/j.neuint.2015.08.013 (2015).
43. Abi-Rached, L., Gilles, A., Shiina, T., Pontarotti, P. & Inoko, H. Evidence of en bloc duplication in vertebrate genomes. *Nature genetics* **31**, doi: 10.1038/ng855 (2002).
44. Ohno, S. *Evolution by gene duplication*. (Springer-Verlag, 1970).
45. Dehal, P. & Boore, J. L. Two rounds of whole genome duplication in the ancestral vertebrate. *PLoS biology* **3**, doi: 10.1371/journal.pbio.0030314 (2005).
46. Brunet, F. G. *et al.* Gene loss and evolutionary rates following whole-genome duplication in teleost fishes. *Molecular biology and evolution* **23**, doi: 10.1093/molbev/msl049 (2006).

47. Kassahn, K. S., Dang, V. T., Wilkins, S. J., Perkins, A. C. & Ragan, M. A. Evolution of gene function and regulatory control after whole-genome duplication: comparative analyses in vertebrates. *Genome research* **19**, doi: 10.1101/gr.086827.108 (2009).
48. Hughes, A. L. The evolution of functionally novel proteins after gene duplication. *Proceedings. Biological sciences / The Royal Society* **256**, doi: 10.1098/rspb.1994.0058 (1994).
49. Force, A. *et al.* Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* **151** (1999).
50. Lynch, M. & Force, A. The probability of duplicate gene preservation by subfunctionalization. *Genetics* **154** (2000).
51. Grone, B. P. & Maruska, K. P. A second corticotropin-releasing hormone gene (CRH2) is conserved across vertebrate classes and expressed in the hindbrain of a basal Neopterygian fish, the spotted gar (*Lepisosteus oculatus*). *The Journal of comparative neurology* **523**, 1125–1143, doi: 10.1002/cne.23729 (2015).
52. Grone, B. P. & Maruska, K. P. Divergent evolution of two corticotropin-releasing hormone (CRH) genes in teleost fishes. *Frontiers in Neuroscience* **9**, doi: 10.3389/fnins.2015.00365 (2015).
53. Chen, C. H. *et al.* Variant GADL1 and response to lithium therapy in bipolar I disorder. *The New England journal of medicine* **370**, 119–128, doi: 10.1056/NEJMoa1212444 (2014).
54. Chatzopoulou, A. *Unraveling the glucocorticoid receptor pathway in zebrafish* PhD thesis, Leiden University, (2012).
55. Lariviere, K., Samia, M., Lister, A., Van Der Kraak, G. & Trudeau, V. L. Sex steroid regulation of brain glutamic acid decarboxylase (GAD) mRNA is season-dependent and sexually dimorphic in the goldfish *Carassius auratus*. *Brain Res Mol Brain Res* **141**, 1–9, doi: 10.1016/j.molbrainres.2005.06.005 (2005).
56. Burki, F. & Kaessmann, H. Birth and adaptive evolution of a hominoid gene that supports high neurotransmitter flux. *Nature Genetics* **36**, 1061–1063, doi: 10.1038/ng1431 (2004).
57. Zhu, J. *et al.* Comparative genomics search for losses of long-established genes on the human lineage. *PLoS computational biology* **3**, e247, doi: 10.1371/journal.pcbi.0030247 (2007).
58. Nishikimi, M., Kawai, T. & Yagi, K. Guinea pigs possess a highly mutated gene for L-gulonon-gamma-lactone oxidase, the key enzyme for L-ascorbic acid biosynthesis missing in this species. *The Journal of biological chemistry* **267**, 21967–21972 (1992).
59. Cui, J., Pan, Y. H., Zhang, Y., Jones, G. & Zhang, S. Progressive pseudogenization: vitamin C synthesis and its loss in bats. *Molecular biology and evolution* **28**, 1025–1031, doi: 10.1093/molbev/msq286 (2011).

Acknowledgements

K.P.M. was supported by startup funds from the College of Science and Department of Biological Sciences at Louisiana State University, a Ralph E. Powe Faculty Enhancement Award from Oak Ridge Associated Universities, and a Louisiana Board of Regents Research Competitiveness Subprogram Grant.

Author Contributions

Both authors had full access to all of the data in this study and take responsibility for its collection and analysis. Study concept and design: B.P.G. and K.P.M. Acquisition of data: B.P.G. Analysis and interpretation of data: B.P.G. and K.P.M. Drafting of the manuscript: B.P.G. Critical revision of the manuscript for important intellectual content: K.P.M. Obtained funding: K.P.M. Administrative, technical, and material support: K.P.M.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Grone, B. P. and Maruska, K. P. Three Distinct Glutamate Decarboxylase Genes in Vertebrates. *Sci. Rep.* **6**, 30507; doi: 10.1038/srep30507 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016