

Effects of Marsh Pond Terracing on Coastal Wintering Waterbirds Before and After Hurricane Rita

Jessica L. O’Connell · John A. Nyman

Received: 2 March 2011 / Accepted: 30 July 2011 / Published online: 28 August 2011
© Springer Science+Business Media, LLC 2011

Abstract From February to March 2005–2006, we surveyed wintering waterbirds to test effects of terracing on coastal pond use before and after Hurricane Rita. Marsh terracing is intended to slow coastal marsh loss in the Chenier Plain by slowing marsh erosion and encouraging vegetation expansion. Terraces also increase marsh edge in ponds, possibly benefiting waterbirds. We monitored paired terraced and unterraced ponds in three sites within southwestern Louisiana’s Chenier Plain. Waterbirds were 75% more numerous in terraced than unterraced ponds. Waterbird richness was similar among ponds when corrected for number of individuals, suggesting terracing increased bird density but did not provide habitat unique from unterraced ponds. Birds were 93% more numerous following Hurricane Rita, mostly due to an influx of migrating waterfowl. Year round residents were similar in number before and after Hurricane Rita. Resident richness did not differ among years after correcting for number of observed individuals. Wading and dabbling foragers were more abundant in terraced ponds and these two guilds represented 74% of birds observed. We detected no difference among ponds for other guilds, i.e., probing, aerial, and diving foragers. Increasing proportion of marsh edge increased bird density disproportionately: On average ponds with 10% edge had 6 birds observed and ponds with 30% edge had 16 birds observed. Terraces increased habitat interspersions and were an effective tool for

increasing numbers of wintering waterfowl and wading birds. The extent to which terraces were sustainable following hurricane forces is unknown.

Keywords Edge effects · Habitat interspersions · Waterfowl · Wading birds · Wetland restoration

Introduction

Louisiana contains 30% of remaining coastal wetlands in the conterminous United States (Field and others 1991) but these wetlands are disappearing rapidly, averaging $16.3 \text{ km}^2 \text{ year}^{-1}$ in the Chenier Plain during 1978–2000 (Barras and others 2003). Louisiana marshes provide important habitat for waterbirds (Keller and others 1984; Myers and others 1987; Martin and Lester 1990; Michot 1996), including wintering and stopover sites for migrants. Continued marsh degradation could therefore affect waterbird population declines locally and regionally. Pond terracing is a novel technique used ubiquitously in Louisiana’s coastal Chenier Plain to slow marsh loss (O’Connell and Nyman 2010). As such, terracing may improve waterbird habitat in the Chenier Plain, with possible benefits for all Central Flyway migrants, but an evaluation of terraced marsh for wintering waterbirds is lacking.

One outcome of marsh loss in the Chenier Plain is interior marsh breakup (Byrnes and others 1995), which may reduce waterbird densities in coastal marsh ponds. Causes of interior marsh breakup in the Chenier Plain are likely varied, but often initiated by human activities. For example, salt-water intrusion along navigation channels causes vegetation die-off in previously fresh or brackish areas, converting emergent marsh into shallow open-water ponds (Turner 1997). Once initiated, wave energy

J. L. O’Connell · J. A. Nyman
School of Renewable Natural Resources, Louisiana State
University Agricultural Center, Baton Rouge, LA 70803, USA
e-mail: jnyman@lsu.edu

J. L. O’Connell (✉)
Department of Zoology, Oklahoma State University,
Stillwater, OK 74078, USA
e-mail: jessica.oconnell@okstate.edu

accelerates erosion along pond edges as they deepen, increasing open-water areas through time (Nyman and others 1994). Lateral marsh erosion possibly is accelerated by global sea level rise and sediment starvation resulting from the channelization of the Mississippi and other rivers (Boesch and others 1994; Turner 1997).

Hurricanes also can cause extensive marsh loss, as occurred when Hurricane Rita, a category 5 storm, struck the Chenier Plain in September 2005, resulting in 217 km² of marsh loss (Barras 2006). Hurricanes cause marsh loss by generating strong wind and waves, churning up marsh areas and over-washing seawater which may kill fresh to brackish vegetation (Chabreck and Palmisano 1973). Hurricanes and other strong storms interact with the Gulf Coast regularly (Gray 1990; Neumann and others 1993). There also is clear evidence of a global increase in tropical storm intensity and duration (Trenberth and others 2007; Elsner and others 2008; Emanuel and others 2008). Such increases may accelerate marsh loss, further impacting Gulf Coast waterbird populations. However, hurricanes and strong storms have complex effects on marshes. While they can cause direct mortality of both vegetation and wildlife, they also are dispersal agents for seeds, vegetative root stock, and nutrient rich off-shore sediments. Therefore, sometimes hurricanes stimulate productivity (Conner and others 1989) and elevation in surviving wetlands (Nyman and others 1995; Turner and others 2006). Some coastal vegetation may be well adapted to storms. For example, Battaglia and Sharitz (2005) suggested storms structured coastal bottomland forest composition and regenerated forests. Waterbirds also may benefit from storm inputs of sediment and nutrients into coastal systems. For example, Convertino and others (2011) demonstrated Snowy Plovers (*Charadrius alexandrinus*) prefer nesting grounds struck the previous season by strong storms, probably because over-washed sediments and storm wrack provided ideal habitat. However, the full effects of hurricanes on waterbirds are not completely known and difficult to test. Consequently, we should evaluate effects of common restoration techniques, such as pond terracing, on waterbird populations in the context of potential marsh instability caused by hurricanes and human-mediated factors.

Pond terracing, ubiquitous in the Chenier Plain, is intended to slow conversion of marsh into open-water ponds by providing a barrier to water movement (Underwood and others 1991; Steyer 1993; Rozas and Minello 2001). Terraces are discontinuous, narrow strips of created marsh formed of dredge material from pond bottoms and stabilized with emergent plants such as *Spartina alterniflora* (Underwood and others 1991; Steyer 1993; Rozas and Minello 2001). Though never evaluated, terraces are thought to reduce wave action in ponds, potentially slowing erosion of pond edges, decreasing water turbidity,

encouraging submerged vegetation production, and decreasing pond depths. Terraced ponds perhaps provide a more hospitable environment for emergent vegetation expansion and increase aerial marsh extent.

Terracing also may support waterbird populations by increasing waterbird forage items, such as nekton, invertebrates, and submerged aquatic vegetation (Rozas and Minello 2001; Whaley and Minello 2002; Barras and others 2003; Bush Thom and others 2004; Cannaday 2006; La Peyre and others 2007; O'Connell and Nyman 2010). These increases likely occur because terracing increases the proportion of marsh edge (boundary between emergent vegetated marsh and open-water) in ponds (O'Connell and Nyman 2010). Marsh edge is a shallow, protected environment which serves as a nursery and refugia for aquatic organisms. Waterbird habitat also may be improved because increasing marsh edge maximizes interspersions of water and vegetated marsh in ponds, i.e. pond cover of open-water and emergent vegetation each approach 50%, as in the hemi-marsh model (Weller and Spatcher 1965). Thus, terracing ponds may improve habitat quality for wintering waterbirds.

We are unaware of previous studies investigating the relationship between wintering waterbirds and habitat interspersions, though this relationship has been demonstrated for breeding waterbirds (Mack and Flake 1980; Kaminski and Prince 1981; Fairbairn and Dinsmore 2001). Additionally, terrace effects on wintering waterbirds have never been evaluated, though during the summer breeding season, we demonstrated that terraces increased waterbird density (O'Connell and Nyman 2010). We originally intended for that previous study to include wintering birds, but the passage of a category 5 storm (Hurricane Rita, 24 September 2005), forced a sampling hiatus and a drastic methodology change. Terrace effects on wintering birds should be evaluated because many Louisiana waterbirds are migrants that only use marsh ponds during winter. These migrants include economically important waterfowl as well as shorebirds. Finally, Hurricane Rita may have caused direct mortality of waterbirds. We have data on wintering waterbird density from 2005, prior to the passage of Hurricane Rita, and from winter 2006, directly following Hurricane Rita. Therefore, we uniquely are able to examine migrating and resident waterbird marsh use in terraced ponds before and after a strong hurricane.

Methodology

Study Area

Our study sites were in coastal southwestern Louisiana's Chenier Plain. This region of the Gulf Coast extends from

west of Vermilion Bay, Louisiana, to High Island, Texas. The Chenier Plain consists of shore-parallel, linear transgressive beach ridges or “cheniers”, separated by broad areas of low-elevation marsh (Penland and Suter 1989). Hurricane Rita struck in September 2005, separating the two wintering periods (February–March 2005 and 2006) we evaluated. Hurricane Rita was an unpredictable event, forcing a change in sampling methods. We minimized these differences by sub-setting our data in both years to equal sampling sites and survey periods. However, some sampling differences remain: 2005 contained one survey in a site not sampled in 2006, plot sizes differed between years (though survey area was approximately equivalent), and time spent in each survey differed. We discuss biases associated with methodological differences and account for them as much as possible.

Site Selection

We monitored pond pairs at each site, one terraced (treatment), and one nearby unterraced pond (control). We selected study ponds by identifying terraced ponds where adjacent marsh was dominated by *Spartina patens*, an indicator of intermediate to brackish conditions. From these, we excluded sites receiving major mineral sediment sources because, unlike Louisiana deltaic wetlands, most Chenier Plain marshes do not receive substantial sediments from rivers, streams or bayous. Unterraced ponds were open-water ponds similar in condition to terraced ponds before terraces were added, i.e. near to the terraced pond and hydrologically connected to it. Study sites were each in different watersheds and hydrologically independent from other sites.

Site Descriptions

Sites differed somewhat in 2005 (before Hurricane Rita) and 2006 (after Hurricane Rita). We monitored waterbirds at two sites each survey period. Ponds averaged 213 ha (range 70–1190 ha). Terraced ponds had more edge habitat (18.3%, SE = 3.1%) than did unterraced ponds (6.4%,

SE = 0.3%), when terraced edge and natural edge were combined into one edge category (Table 1). Sites used were in Sabine National Wildlife Refuge (NWR) (terraced pond coordinates: 0434212 East and 3304478 North, Unit 5 and unterraced pond coordinates: 0434609 East and 3311970 North, Unit 6, hydraulically connected to Unit 5). Sweet Lake (coordinates: 480933 East and 3312269 North, owned by Sweet Lake Gas and Oil Co. and Miami Corporation), and an impounded unit in Rockefeller State Wildlife Refuge (SWR) (coordinates: 523597 East and 3284941 North, Unit 4, owned by Louisiana Department of Wildlife and Fisheries). Coordinates are UTM NAD 1984, Zone 15 N. Terraces were constructed in Sweet Lake in 2001, in Rockefeller SWR in 2002, and in Sabine in 2001. Waterfowl hunting occurred in Sweet Lake during January months in terraced and unterraced ponds equally. Hunting was not allowed in Rockefeller SWR or Sabine NWR. The unterraced pond at Sabine NWR was terraced after our first survey in 2005. We discontinued sampling this site and replaced it with the Sweet Lake site. We retained Sabine NWR in our study so that two sites are sampled in all surveys in both years.

Survey Methods 2005 (Pre-Hurricane)

Surveys were conducted every two-weeks from 11 February 2005 through 20 March 2005. Each pond contained multiple survey plots. One plot/pond was sampled each survey session; different plots were sampled in subsequent sessions. Ultimately, sampling effort was even among plots. Survey methods are as described in O’Connell and Nyman (2010), but briefly, plots were delineated by marking plot corners with PVC pipe. Plots ranged from 9–12 ha (average 11 ha) due to geographical constraints. Observers were dropped off at plots by boat. One observer counted plots while hidden in emergent vegetation. Terraced and unterraced ponds were surveyed simultaneously by separate observers. Surveys began at dawn. To minimize double-counting, observers counted birds by scanning plots from end to end, disregarding birds that entered or left plots after an area had been counted. Observers counted

Table 1 Area (ha) of water classified as open-water (OW), water near a natural edge (NE), terraced edge (TE), total water near any edge (Total E), the ratio of TE:NE, and the percent of water that is edge habitat

Site	Pond type	OW	TE	NE	Total E	TE:NE ratio	% edge
Sabine	Terraced	427.8	35.5	9.8	45.3	1.99	9.6
	Unterraced	1099.7	–	91.1	91.1	–	7.6
Sweet Lake	Terraced	155.5	50.5	10.3	60.8	4.9	28.1
	Unterraced	95.5	–	5.4	5.4	–	5.4
Rockefeller	Terraced	59.3	7.3	5.1	12.3	1.4	17.2
	Unterraced	72.3	–	4.8	4.8	–	6.2

birds after a 15 min settling period. Observers counted plots multiple times, but for comparison with 2006 survey methods, only the first plot count was used in analyses.

Survey Methods 2006 (Post-Hurricane)

Field housing was unavailable in the Chenier Plain following Hurricane Rita. We therefore designed surveys to maximize ponds visited during a single day and surveyed all plots/pond each survey session. To minimize bias associated with time of day, surveys began at different ponds and plots each survey session, such that sampling effort was even among ponds. Hurricane wreckage prevented access to the Chenier Plain from September to early November 2005. During December, we tested survey methods. We inferred from these trials that boat noise disturbance to birds was reduced when plots were approached at low speeds and oblique angles, boats stopped at a distance from birds, and small discrete plots were counted rather than large plots or long, narrow transects. Flushing of birds was minimal using these methods, though some did occur. However, our view was unobstructed because ponds are large and open, and we successfully determined bird locations prior to boat disturbance. Thus, we ensured we counted birds using plots and ignored birds flushed from adjacent marsh.

Plots were delineated with PVC as in 2005. Plot locations overlapped 2005 plots, but plot size was reduced from 10 ha average, allowing quick, complete bird counts. Plots ranged from 3–5 ha (average 4.3 ha) due to geographical constraints. We surveyed 2–3 plots in ponds during each survey rather than one plot as was done in 2005. Total survey area in ponds was approximately equivalent to plot size surveyed in 2005.

Two observers and one boat driver conducted surveys. Plots were counted upon arrival via boat, rather than after a 15 min settling period as was done in 2005. We interpret this bias in our discussion. Each observer independently recorded observations and the same observers counted both

terraced and unterraced ponds. We used the average number for each species counted by both observers as our species abundance estimate. We sampled sites once every two weeks from 7 February 2006 to 29 March 2006, generating equal survey periods between years.

Statistical Methods

We grouped bird species into foraging guilds to compare numbers of individuals between pond types. We used foraging classifications of De Graaf and others (1985), except that we classified Common Moorhen as dabblers based on our behavior observations. Guilds were as follows: diving foragers, dabbling foragers, wading foragers, probing foragers, and aerial foragers (Table 2). We also grouped year round residents to compare whether species present during Hurricane Rita's passage differed in number among years (Table 3). We compared number of total birds, divers, dabblers, waders, probers, aerialists, and year round residents with general linear mixed models using Poisson error (lmer program using R) and random intercepts by site and plot size to account for correlated repeated measurements within sites and variation due to plot size (Zuur and others 2009). We reduced independent variables in models if explanatory variables were not significant ($P > 0.1$). We used EstimateS (Colwell 2005) to generate species accumulation curves, accounting for rarefaction in species richness among treatments (Pielou 1975; Palmer and others 2008). We used 500 randomizations and plot moment-based estimation (Mao Tau estimator in EstimateS) of species richness versus number of individuals as recommended by Colwell and others (2004).

Additionally, we used a general linear mixed model with Poisson error and random site and plot size to compare number of all birds observed with proportion of pond within 10 m of marsh edge (e.g., habitat interspersion). We used ArcGIS 9.1 (ESRI Corp, Redlands, CA) to compute the proportion of open-water and water near marsh edge in each pond type by analyzing 2004 DOQQ aerial

Table 2 General linear models of the relationship among bird density (birds ha⁻¹ ± SE), pond type (factor), survey week (continuous), and year (factor), with Poisson error structure and log-link between the model and response

Model	Intercept	Pond type (unterraced)	Year (2006)	Survey week
Total bird density	-1.18 ± 0.82*	-1.17 ± 0.46***	2.95 ± 0.82***	0.45 ± 0.05***
Dabbler density	-4.78 ± 1.57**	-1.8 ± 0.84*	4.33 ± 1.47**	0.95 ± 0.09***
Wader density	-1.61 ± 0.72*	-1.39 ± 0.65*	1.89 ± 0.76*	NS
Prober density	NS	NS	NS	NS
Aerial density	NS	NS	NS	NS
Diver density	NS	NS	NS	NS
Density of residents	1.12 ± 0.61~	-1.15 ± 0.57*	NS	NS

*** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$; and ~ $P < 0.1$

Table 3 Common and scientific names and mean number of individuals (\pm SE) for observed species and foraging guilds

Species	2005		2006	
	Terraced	Unterraced	Terraced	Unterraced
Diving foragers	0.1 \pm 0.33	0.1 \pm 0.17	0.3 \pm 0.21	0.2 \pm 0.79
^Pied-billed Grebe (<i>Podilymbus podiceps</i>)	0.1 \pm 0.13	0.3 \pm 0.25	0.3 \pm 0.25	0.3 \pm 0.16
Common Loon (<i>Gavia immer</i>)	0 \pm 0	0 \pm 0	0 \pm 0	0.3 \pm 0.16
^Double-crested Cormorant (<i>Phalacrocorax auritus</i>)	0.9 \pm 0.58	0 \pm 0	0.4 \pm 0.38	3.0 \pm 3.00
Wading foragers	0.3 \pm 0.21	0.1 \pm 0.3	1.3 \pm 0.8	0.3 \pm 0.21
^Great Blue Heron (<i>Ardea herodias</i>)	0.1 \pm 0.13	0 \pm 0	0.4 \pm 0.26	0 \pm 0
^Tricolored Heron (<i>Egretta tricolor</i>)	0 \pm 0	0 \pm 0	0.1 \pm 0.13	0 \pm 0
^Great Egret (<i>Ardea alba</i>)	0.1 \pm 0.13	0 \pm 0	0.8 \pm 0.31	0.3 \pm 0.16
^Snowy Egret (<i>Egretta thula</i>)	0.1 \pm 0.13	0 \pm 0	0 \pm 0	0 \pm 0
^White Ibis (<i>Eudocimus albus</i>)	0 \pm 0	0 \pm 0	0.1 \pm 0.13	0 \pm 0
Dabbling foragers	1.4 \pm 0.31	0.3 \pm 0.05	29.1 \pm 9.46	5.4 \pm 1.92
Blue-winged Teal (<i>Anas discors</i>)	0 \pm 0	0 \pm 0	5.9 \pm 2.80	2.4 \pm 2.38
Gadwall (<i>Anas strepera</i>)	1.3 \pm 0.73	0.3 \pm 0.25	5.1 \pm 4.29	1 \pm 0.42
Green-winged Teal (<i>Anas crecca</i>)	0 \pm 0	0 \pm 0	26.0 \pm 12.97	0.9 \pm 0.88
Mallard (<i>Anas platyrhynchos</i>)	0 \pm 0	0 \pm 0	0.1 \pm 0.13	0 \pm 0
^Mottled Duck (<i>Anas fulvigula</i>)	0 \pm 0	0.3 \pm 0.25	0.3 \pm 0.25	0.6 \pm 0.42
Northern Pintail (<i>Anas acuta</i>)	0 \pm 0	0 \pm 0	0.5 \pm 0.5	0 \pm 0
Northern Shoveler (<i>Anas clypeata</i>)	0 \pm 0	0 \pm 0	5.1 \pm 2.29	2.5 \pm 2.50
^Common Moorhen (<i>Gallinula chloropus</i>)	0 \pm 0	0.1 \pm 0.13	0 \pm 0	0 \pm 0
Probing foragers	0 \pm 0	0 \pm 0	0 \pm 0	0.3 \pm 0.33
^Black-necked Stilt (<i>Himantopus mexicanus</i>)	0 \pm 0	0 \pm 0	0.1 \pm 0.13	0 \pm 0
Greater Yellowlegs (<i>Tringa melanoleuca</i>)	0 \pm 0	0 \pm 0	0.5 \pm 0.5	0.6 \pm 0.63
Lesser Yellowlegs (<i>Tringa flavipes</i>)	0 \pm 0	0 \pm 0	0 \pm 0	0.4 \pm 0.38
^Willet (<i>Catoptrophorus semipalmatus</i>)	0 \pm 0	0 \pm 0	1.5 \pm 1.5	0 \pm 0
Aerial foragers	0.7 \pm 0.21	0.2 \pm 0.17	0.7 \pm 0.21	1.0 \pm 0.45
^Laughing Gull (<i>Larus atricilla</i>)	0 \pm 0	0.1 \pm 0.13	0.6 \pm 0.63	0 \pm 0
Ring-billed Gull (<i>Larus delawarensis</i>)	0 \pm 0	0 \pm 0	0.6 \pm 0.38	0.3 \pm 0.16
^Caspian Tern (<i>Sterna caspia</i>)	0.3 \pm 0.16	0 \pm 0	0.6 \pm 0.38	0 \pm 0
^Forster's Tern (<i>Sterna forsteri</i>)	0.4 \pm 0.38	0.4 \pm 0.38	2.4 \pm 1.46	1.3 \pm 1.25
^Royal Tern (<i>Sterna maxima</i>)	0 \pm 0	0 \pm 0	0.8 \pm 0.62	0 \pm 0
Year round residents	1.0 \pm 0.48	0.3 \pm 0.33	5.0 \pm 2.58	1.6 \pm 1
Total birds	2.9 \pm 1.25	0.7 \pm 0.25	38.1 \pm 9.97	9.6 \pm 2.11

^ Year round residents

photographs of sites (See O'Connell and Nyman 2010 for methods).

Results

Interaction terms between pond type, survey week and year were not significant for any response variable. Birds were 75% more numerous in terraced ponds than unterraced (Fig. 1). When species richness was corrected for number of individuals (rarefaction), richness was similar among pond types, i.e. confidence intervals for species accumulation curves overlap though the number of individuals was

not equal (Fig. 2a, b). Birds were 93% more numerous following Hurricane Rita, mostly due to influx of migrating dabbling ducks (Fig. 1). Numbers of resident birds was roughly equivalent between years (Table 3). Resident species richness also was similar among years when numbers of individuals were accounted for (Fig. 2c).

After combining pond types and years guilds were, in order of lowest average density to highest: divers (0.2, SE = 0.4), probers (0.3, SE = 0.3), waders (0.5, SE = 0.04), aerialists (1.1, SE = 0.7), and dabblers (9.0, SE = 2.7).

Dabbling birds and wading birds were 82 and 77% more numerous in terraced ponds than in unterraced ponds




Fig. 1 Number of waterbird foraging guilds and resident species in terraced and unterraced ponds (mean ± SE), in winter of 2005 and 2006, Chenier Plain, Louisiana. Hurricane Rita struck between years

(Table 3; Fig. 1) and together comprised 74% of all birds observed. Dabblers and waders were 95 and 77% more numerous in 2006 than 2005, respectively. Aerialists were somewhat abundant (Fig. 1), but we did not detect a difference in numbers among pond types or years. Probing and diving foragers were not abundant, and we also did not detect differences among pond types or years. Resident species were 80% more numerous in terraced ponds than unterraced, but we detected no difference in numbers between years (Table 3; Fig. 1). Finally, there were greater number of birds in ponds with greater proportion of marsh edge (Model: total number of birds = $1.3 + 4.9 \times$ (proportion edge), Poisson error structure and log link between response and model; Fig. 3).

Discussion

Terrace Effects

Waterbirds were more numerous in terraced ponds, probably because terracing increased the proportion of marsh edge (O’Connell and Nyman 2010) and enhanced habitat




Fig. 2 Species accumulation curves of all species among pond types in 2005 (a), 2006 (b), and resident species among years (c), depicting moment-based estimation of observed species richness versus number of individual birds in ponds




Fig. 3 Number of birds explained by proportion of marsh edge in ponds in winter 2005 and 2006 Chenier Plain, LA (Model: total birds = $1.3 + 4.9 \times$ (proportion of edge), Poisson error structure and log-link between response and model)

complexity (Feagin and Wu 2006). Birds likely were also more numerous in ponds with greater marsh edge because this increases habitat interspersions of cover and open-water. Ours is the first demonstration of a habitat interspersions effect on wintering waterbirds. However, we demonstrated similar effects in these sites for spring and summer waterbirds during 2005 (O’Connell and Nyman 2010). Others have shown the importance of habitat interspersions for breeding waterbirds in northern marshes (Weller and Spatcher 1965; Mack and Flake 1980; Kaminski and Prince 1981; Fairbairn and Dinsmore 2001). Additionally habitat interspersions has other ecological influences. For example, habitat interspersions has increased macroinvertebrate densities, improved water quality, reduced mosquito populations, and increased mosquito predator populations (Thullen and others 2002, 2005). Therefore, maximizing habitat interspersions in marshes may improve multiple ecological services, potentially benefiting waterbirds.

Foraging guilds varied in their response to pond terracing. Waders and dabblers were more numerous in terraced ponds. Waterbirds that forage either for SAV or for items dependent on SAV for food, habitat, or refuge likely benefited from terracing because of increased SAV (Cannaday 2006; O’Connell and Nyman 2010), invertebrates (Whaley and Minello 2002), and nekton (Minello and Rozas 2002; La Peyre and others 2007; O’Connell and Nyman 2010) adjacent to terraced edges. Terracing should have therefore improved densities of SAV-foraging dabblers and piscivorous waders.

We did not detect differences among pond types for aerialists, divers and probers. During spring and summer, aerial foragers in these sites used terraced ponds more often (O’Connell and Nyman 2010). Aerialists may behave differently during winter than during the breeding season, when proximity to nest sites and provisioning of young are important. For example, Ring-billed Gulls have been frequently observed foraging along terrestrial edges during summer (Baird 1976; Welham 1987), but often observed plunge diving for fish in shallow open-water during winter (Chapman and Parker 1985). Diving foragers were not abundant and did not differ in their use of ponds in either winter (this study) or summer (O’Connell and Nyman 2010). Diving foragers likely do not rely on shallow ponds as important habitat (ponds studied were less than 1 m deep). They also may feed on forage items from pond bottoms, rather than those close to pond edges. Probing foragers also were not abundant and did not differ among pond types over winter. Probers such as shorebirds may not have been numerous because winter water depths were too deep. Water depths possibly were deeper because of undrained storm surge following Hurricane Rita (see discussion below). However, few shorebirds winter in

Louisiana and generally are more abundant during spring and fall migration. Terraced ponds could be improved as shorebird habitat if terraces had broader widths and gentler slopes, increasing accessible shallow areas.

It is possible unterraced ponds provide habitat characteristics not present in terraced ponds. However, there is little evidence to support this. For example, species richness was similar among pond types when numbers of individuals were accounted for. Conclusive evidence for habitat heterogeneity among ponds would be provided only if species richness were independent of density (Fridley and others 2006).

Hurricane Effects

Hurricane Rita hit the Chenier Plain in September 2005, causing extensive destruction of coastal development and substantial upheaval in adjacent marshes. Numerous “marsh balls” of uprooted *S. patens* were deposited in previously open-water areas of our study ponds. Undrained storm surge remained in ponds, likely elevating salinities. Fish kills were observed on adjacent uplands, perhaps reducing food sources. However, the hurricane also may have brought in additional nutrients and food sources from offshore areas. We observed no alteration of terrace widths or of surrounding natural marsh edge morphology. The sides of terrace edges also did not appear undercut directly following the hurricane. However, we revisited our study sites in Rockefeller SWR in late spring 2006 following the completion of this study. Terraces at this impounded site were vastly degraded, perhaps because sustained high salinity killed vegetation. Effects of hurricanes on terraced marsh are undetermined. If constant repair of eroding terraces is necessary, many marsh functions may never return to pre-disturbance conditions. Long-term monitoring is necessary to determine if terraces are sustainable.

Hurricane Rita did not appear to reduce bird numbers during this study. Species observed were typical for winter in coastal Louisiana. Many dabbling ducks and shorebirds were migrants not present during the hurricane’s passage. Total waterbird numbers observed in plots were greater following the hurricane than during the previous winter, or than observed during summer at these same sites (O’Connell and Nyman 2010). Higher numbers particularly were observed for migrating dabbling ducks following Hurricane Rita. There is little reason to presume the hurricane directly influenced migrant density. One explanation for higher numbers could be plot size differences among years. Smaller plots, such as those sampled following the hurricane, are known to inflate density estimates over estimates from larger plots because of larger plot edge to plot area ratios (White and others 1982). We minimized this bias by modeling individuals rather than density and

Table 4 Temperature and rainfall in southwestern Louisiana during February and March of 2005 and 2006

Year	Month	Average temperature	Departure from normal	Total rainfall	Departure from normal
2005	January	55	4	3.17	-3.17
2005	February	56	1	7.55	3.59
2005	March	59	-2	2.69	-1.74
2006	January	57	6	2.7	-3.34
2006	February	53	-2	4.54	0.58
2006	March	64	3	1.22	-3.21

Data from the Louisiana Office of State Climatology (<http://www.losc.lsu.edu/>)

including plot area as a covariate in analyses. However, this well known sampling artifact may still account for part of observed between year differences. Further, locally, weather was warmer and drier in 2006 than 2005 (Table 4). However, continent-wide weather patterns or region-wide hunting pressure were likely as influential as local conditions. Sampling during 2005 was amidst El Nino conditions, while 2006 sampling was at the end of La Nina conditions (NOAA 2011). La Nina sometimes has been linked with severe droughts in North American northern prairies (Trenberth and Guillemot 1996), where most waterfowl migrants breed. Corroborating this, a record drought persisted during summer 2005 across American mid-west prairies (Palecki and Hilberg 2006), perhaps limiting grain foods and encouraging waterfowl migration southwards. Year round residents did not differ among years, implying either that there was not substantial direct hurricane mortality or that immigration offset mortality. Only two species were observed in 2005 and not 2006: Common Moorhen and Snowy Egret. Other resident species were either roughly equivalent between years or were more numerous following Hurricane Rita. Leberg and others (2007) found in their comparison of nesting colonial waterbirds that nesting pairs for most species increased following Hurricanes Rita and Katrina in southern Louisiana. They also documented substantial movement of birds between colony sites. One further study has documented hurricane effects on waterbird communities, in this case following Hurricane Hugo in South Carolina (Shepherd and others 1991). They documented increases for some species in study sites following the hurricane while others vastly declined, but could not definitively attribute declines to mortality or emigration. Studies of non-waterbird species also have documented bird declines in some habitats and increases in others and generally have ascribed this to populations tracking dispersed food and resources following hurricanes (Wunderle and others 1992 and references therein). Generally, resident waterbirds fair better if hurricanes strike outside the breeding season. Brackish species also are more resilient than beach and fresh marsh species because of high wave and salt-water impacts (Michener and others 1997). Taken altogether, we suggest

Hurricane Rita caused a redistribution of resources rather than substantial waterbird mortality in these sites; however, evidence is not conclusive.

Sampling differences between seasons could have influenced our results, but likely did not. During winter 2005 we counted larger plots. Following the hurricane, we counted smaller plots, which has often been associated with finding fewer individuals and species (Rosenzweig 1995) and especially fewer rare species (Tjørve and others 2008), though it also inflates density estimates (White and others 1982), as discussed above. We minimized area bias by adding multiple small plots together to achieve roughly equivalent survey areas between years and modeling plot size bias in our analyses. Additionally, following the hurricane, surveys were conducted via boat rather than while hidden in emergent vegetation and settling periods prior to counting differed. We therefore may have counted fewer secretive birds in 2006. Despite this bias toward detecting fewer species following the hurricane, species richness was equivalent between years, further implying that direct hurricane mortality was minimal.

Summary and Conclusions

Overall, terraces increased the proportion of edge in ponds and numbers of wintering birds by 93%, a dramatic increase. Terraces benefited waterbirds by increasing habitat interspersion in ponds, generating ideal waterbird conditions. Further improvements in waterbird habitat may be achieved by building terraces with shallower slopes, which could increase water depth variation in ponds and promote use by shorebirds. We do not know whether Hurricane Rita caused significant bird mortality, but for remaining birds, terracing increased their use of coastal marsh ponds. Further monitoring is necessary to determine if use of pond terracing is sustainable both within and outside the Chenier Plain. However, habitat interspersion benefits waterbirds in both winter and summer.

Acknowledgments This project was funded by USFWS Gulf Coast Joint Venture and Ducks Unlimited Inc., project number LA-86-1.

We thank Rockefeller State Wildlife Refuge, Vermilion Corporation, Sweet Lake Gas and Oil Co., and Miami Corporation for allowing site access.

References

- Baird PA (1976) Comparative ecology of California and ring-billed gulls (*Larus californicus* and *L. delawarensis*). PhD dissertation, University of Montana
- Barras JA (2006) Land area changes in coastal Louisiana after the 2005 hurricanes: a series of three maps. United States Geological Survey, Washington
- Barras JA et al (2003) Historical and projected coastal Louisiana land changes: 1978–2050. United States Geological Survey, Washington, Open File Report 03-334
- Battaglia LL, Sharitz RR (2005) Effects of natural disturbance on bottomland hardwood regeneration. In: Fredrickson LH, King SL, Kaminski RM (eds) Ecology and Management of bottomland hardwood systems: the state of our understanding. Gaylord Memorial Laboratory, University of Missouri-Columbia, pp 121–136
- Boesch DF, Josselyn MN, Mehta AJ, Morris JT, Nuttle WR, Simenstad CA, Swift DJP (1994) Scientific assessment of coastal wetland loss, restoration, and management in Louisiana. Journal of Coastal Research Special Issue 20:1–103
- Bush Thom CS, La Peyre MKG, Nyman JA (2004) Evaluation of nekton use and habitat characteristics of restored Louisiana marsh. Ecological Engineering 23:63–75
- Byrnes MR, McBride RA, Tao Q, Duvic L (1995) Historical shoreline dynamics along the Chenier Plain of southwestern Louisiana. Gulf Coast Association of Geological Societies Transactions XLV:113–122
- Cannaday CD (2006) Effects of terraces on submerged aquatic vegetation in shallow marsh ponds in coastal southwestern Louisiana. MS Thesis, Louisiana State University
- Chabreck RH, Palmisano AW (1973) The effects of Hurricane Camille on the marshes of the Mississippi River Delta. Ecology 54:1118–1123
- Chapman BA, Parker JW (1985) Foraging areas, techniques, and schedules of wintering gulls on southeastern Lake Erie. Colonial Waterbirds 8:135–141
- Colwell RK (2005) EstimateS: statistical estimation of species richness and shared species from samples. Version 8.2. R. Colwell, <http://purl.oclc.org/estimates>
- Colwell RK, Mao CX, Chang J (2004) Interpolating, extrapolating, and comparing incidence-based species accumulation curves. Ecology 85:2717–2727
- Conner W, Day J, Baumann R, Randall J (1989) Influence of hurricanes on coastal ecosystems along the northern Gulf of Mexico. Wetlands Ecology and Management 1:45–56
- Convertino M, Elsner JB, Muñoz-Carpena R, Kiker GA, Martinez CJ, Fischer RA, Linkov I (2011) Do tropical cyclones shape shorebird habitat patterns? Biogeoclimatology of snowy plovers in Florida. PLoS One 6:e15683
- De Graaf RM, Tilghman NG, Anderson SH (1985) Foraging guilds of North American birds. Environmental Management 9:493–536
- Elsner JB, Kossin JP, Jagger TH (2008) The increasing intensity of the strongest tropical cyclones. Nature 455:92–95
- Emanuel K, Sundararajan R, Williams J (2008) Hurricanes and global warming: results from downscaling IPCC AR4 simulations. Bulletin of the American Meteorological Society 89:347–367
- Fairbairn SE, Dinsmore JJ (2001) Local and landscape-level influences on wetland bird communities of the prairie pothole region of Iowa, USA. Wetlands 21:41–47
- Feagin RA, Wu XB (2006) Spatial pattern and edge characteristics in restored terrace versus reference salt marshes in Galveston Bay. Wetlands 26:1004–1011
- Field D, Reyer A, Genovese P, Shearer B (1991) Coastal wetlands of the United States: an accounting of a valuable national resource. National Oceanic and Atmospheric Administration, Rockville
- Fridley Jason D, Peet Robert K, van der Maarel E, Willems Jo H (2006) Integration of local and regional species-area relationships from space-time species accumulation. The American Naturalist 168:133–143
- Gray WM (1990) Strong association between West African rainfall and U.S. landfall of intense hurricanes. Science 249:1251–1256
- Kaminski RM, Prince HH (1981) Dabbling duck and aquatic macroinvertebrate responses to manipulated wetland habitat. Journal of Wildlife Management 45:1–15
- Keller CE, Spendelow JP, Greer RD (1984) Atlas of wading bird and seabird colonies in coastal Louisiana, Mississippi, and Alabama: 1983. U. S. Fish and Wildlife Service, Washington, FWS/OBS-84/13
- La Peyre MK, Gossman B, Nyman JA (2007) Assessing functional equivalency of nekton habitat in enhanced habitats: comparison of terraced and unterraced marsh ponds. Estuaries and Coasts 30:526–536
- Leberg PL, Green MC, Adams BA, Purcell KM, Luent MC (2007) Response of waterbird colonies in southern Louisiana to recent drought and hurricanes. Animal Conservation 10:502–508
- Mack GD, Flake LD (1980) Habitat relationships of waterfowl broods on South Dakota stock ponds. Journal of Wildlife Management 44:695–700
- Martin RP, Lester GD (1990) Atlas and census of wading bird and seabird nesting colonies in Louisiana: 1990. Louisiana Department of Wildlife and Fisheries, Natural Heritage Program, Baton Rouge, LA, Special Publication No. 3
- Michener WK, Blood ER, Bildstein KL, Brinson MM, Gardner LR (1997) Climate change, hurricanes and tropical storms, and rising sea level in coastal wetlands. Ecological Applications 7:770–801
- Michot TC (1996) Marsh loss in coastal Louisiana: implications for management of North American Anatidae. Gibier Faune Sauvage Game and Wildlife 13:941–957
- Minello TJ, Rozas LP (2002) Nekton in Gulf Coast wetlands: fine-scale distributions, landscape patterns, and restoration implications. Ecological Applications 12:441–455
- Myers J, Morrison R, Antas P, Harrington B, Lovejoy T, Sallaberry M, Senner S, Tarak A (1987) Conservation strategy for migratory species. American Scientist 75:18–26
- Neumann CJ, Jarvinen BR, McAdie CJ, Elms JD (1993) Tropical cyclones of the North Atlantic Ocean, 1871–1992. National Climatic Data Center, Asheville
- NOAA (2011) Multivariate ENSO Index. <http://www.esrl.noaa.gov/psd/enso/mei/mei.html>. Accessed March 1, 2011
- Nyman JA, Carlross M, DeLaune RD, Patrick WH (1994) Erosion rather than plant dieback as the mechanism of marsh loss in an estuarine marsh. Earth Surface Processes and Landforms 19:69–84
- Nyman JA, Crozier CR, DeLaune RD (1995) Roles and patterns of hurricane sedimentation in an estuarine marsh landscape. Estuarine, Coastal and Shelf Science 40:665–679
- O’Connell JL, Nyman JA (2010) Marsh terraces in coastal Louisiana increase marsh edge and densities of waterbirds. Wetlands 30:125–135
- Palecki M, Hilberg SD (2006) Regional analysis of the 2005 drought. In: Kunkel KE (ed) The 2005 Illinois drought. Illinois State Water Survey, Champaign, pp 26–37
- Palmer MW, McGlenn DJ, Fridley JD (2008) Artifacts and artificions in biodiversity research. Folia Geobot 43:245–257

- Penland S, Suter JR (1989) The geomorphology of the Mississippi River Chenier Plain. *Marine Geology* 90:231–258
- Pielou EC (1975) *Ecological diversity*. Wiley, New York
- Rosenzweig ML (1995) *Species diversity in space and time*. Cambridge University Press, Cambridge
- Rozas LP, Minello TJ (2001) Marsh terracing as a wetland restoration tool for creating fishery habitat. *Wetlands* 21:327–341
- Shepherd P, Crockett T, Santo TLD, Bildstein KL (1991) The impact of Hurricane Hugo on the breeding ecology of wading birds at Pumpkinseed Island, Hobcaw Barony, South Carolina. *Colonial Waterbirds* 14:150–157
- Steyer GD (1993) Final report Sabine terracing project. Louisiana Department of Natural Resources, Coastal Restoration Division, Baton Rouge
- Thullen JS, Sartoris JJ, Walton WE (2002) Effects of vegetation management in constructed wetland treatment cells on water quality and mosquito production. *Ecological Engineering* 18:441–457
- Thullen JS, Sartoris JJ, Nelson SM (2005) Managing vegetation in surface-flow wastewater-treatment wetlands for optimal treatment performance. *Ecological Engineering* 25:583–593
- Tjørve E, Kunin WE, Polce C, Calf Tjørve KM (2008) Species–area relationship: separating the effects of species abundance and spatial distribution. *Journal of Ecology* 96:1141–1151
- Trenberth KE, Guillemot CJ (1996) Physical processes involved in the 1988 drought and 1993 floods in North America. *Journal of Climate* 9:1288–1298
- Trenberth KE et al (2007) Observations: surface and atmospheric climate change. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Millers HL (eds) *Climate change 2007 : the physical science basis : contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp 747–845
- Turner RE (1997) Wetland loss in the northern Gulf of Mexico: multiple working hypotheses. *Estuaries* 20:1–13
- Turner RE, Baustian JJ, Swenson EM, Spicer JS (2006) Wetland sedimentation from Hurricanes Katrina and Rita. *Science* 314:449–452
- Underwood SG, Steyer GD, Good B, Chambers D (1991) Bay bottom terracing and vegetative planting: an innovative approach for habitat and water quality enhancement. In: Webb FJ (ed) *Annual Conference on Wetlands Restoration and Creation*. Tampa, FL, pp 164–173
- Welham CVJ (1987) Diet and foraging behavior of ring-billed gulls breeding at Dog Lake, Manitoba. *The Wilson Bulletin* 99: 233–239
- Weller MW, Spatcher CS (1965) Role of habitat in the distribution and abundance of marsh birds. Agricultural and Home Economic Experiment Station, Iowa State University of Science and Technology, Ames
- Whaley SD, Minello TJ (2002) The distribution of benthic infauna of a Texas salt marsh in relation to the marsh edge. *Wetlands* 22:753–766
- White GC, Anderson DR, Burnham KP, Otis DL (1982) Density estimation. Capture-recapture and removal methods for sampling closed populations. Los Alamos National Laboratory, Los Alamos, pp 120–131
- Wunderle JM Jr, Lodge DJ, Waide RB (1992) Short-term effects of Hurricane Gilbert on terrestrial bird populations on Jamaica. *The Auk* 109:148–166
- Zuur A, Ieno E, Walker N, Saveliev A, Smith G (2009) *Mixed effects models and extensions in ecology with R*. Springer, New York